

# How many acres of potatoes does a society need?

N T Moore

‡

E-mail: nmoore@winona.edu

**Abstract.** One of the main difficulties in a class on Sources of Energy and Social Policy is the wide variety of units used by different technologists (BTU's, Barrels of oil, Quads, kWh, etc). As every student eats, I think some of this confusion can be resolved by starting and grounding the class with a discussion of food and food production. A general outline for this introduction is provided and two interesting historical cultural examples, Tenochtitlan and the Irish Potato Famine, are provided. Science and Social Policy classes are full of bespoke units and involve many different contexts. Starting the class with a discussion of food energy is a nice way for everyone to start with the same context. In addition, discussion of Food Energy can lead to interesting historical claims.

*Keywords:* Energy, Social Policy, kcals, Tenochtitlan, Irish Potato Famine, History, self-reliance Submitted to: *Phys. Educ.*

‡ Present address: Department of Physics, Winona State University, Winona, MN 55987, USA

## 1. Introduction

When the United States entered World War One one of the problems they faced was logistics. How much food do you need to ship overseas to Europe to feed a million soldiers? That early work in nutrition led to the 3000 Calorie diet many people remember from secondary Health Education class. A bit about units you might remember: 1 *Calorie* = 1 *kilo – calorie (kcal)*, and a dietician might build a 3000 kcal diet for a 20 year old basketball player. A *calorie* is the amount of energy it takes to heat a gram of water by a degree Celsius. There are about 4.2 Joules in a single calorie, and a Joule occurs all over introductory physics. If you need to buy a new home furnace, the sales brochure might advertise that it is capable of delivering 100,000 BTU's of heat each hour. What's a BTU? Heat a pound of water by  $1^{\circ}F$ . Of course Heat Pumps are far more efficient than simply burning methane or propane, but they consume kilowatt-hours (kWh) of electricity, not BTU's. What's a kWh? Run a 1000 Watt toaster for an hour and you'll have pulled one kWh off the grid, it will cost you about \$0.13 in Minnesota. If you decide to put solar panels in your backyard, they will probably collect about 10% of the 3.5kWh the sun delivers to each square meter of your lawn (in Minnesota) each day.

As the previous paragraph illustrates, there are a frustratingly large number of different units in an "Energy" class. At Winona State, this 3 credit class fulfills a "Science and Social Policy" general education requirement and is taken by students from across the university. Lots of college majors don't require a math class beyond algebra or introductory statistics and the population is largely math-averse. You could jokingly say that one of the main things students learn in the class is unit conversion, but it isn't far off. Nearly every field finds energy a useful representation, and every profession has their own set of units and terminology that's most well suited for quick calculation. Would a medical lab scientist talk about the fractional acre-foot of urine needed test kidney function? No, but someone in the central valley of California would certainly care about the acre-feet of water necessary to grow almonds! Does a gas station price their gasoline in dollars per kWh? Given the growing electrification of cars, they might soon.

Everyone eats, maybe not 3000 kcals per day, but at least something every day. When I teach our energy class, I spend a few weeks talking about food energy before all other types. While food production is not central to climate change and wars over oil, food is essential in a way that diesel and gasoline are not. Vehicle fuel makes modern life possible, but we could live, unpleasantly, without it. We can't live without fats and protein.

## 2. Food Energy

To introduce Food Energy, I ask the students to work through a few questions:

## 2.1. food converted into body heat

Planning to save money, one college student decides to go to an all-you-can-eat buffet each day at 11am. If he brings homework and stretches the meal out for a few hours he can get all 3000 *kcal*s with only one meal bill. Food is fuel for the human body – could too much fuel make his body feel sick? If his body burned all this food at once, how much warmer would he get? Useful information: the student has a mass of 80kg and is made mostly of water. A Calorie heats 1 kg of water 1°C.

Possible Answer: We'll equate food energy with calorimetric heating and assume human bodies have the same heat capacity as water, about  $1\text{kcal}/\text{kg} \cdot \text{C}^\circ$ . This allows us to calculate the body's temperature increase.

$$3000\text{kcal} = 80\text{kg} \cdot 1 \frac{\text{kcal}}{\text{kg} \cdot \text{C}^\circ} \cdot \Delta T \quad (1)$$

$$\Delta T \approx + 37.5\text{C}^\circ \quad (2)$$

Students are normally quite surprised at this answer. Although it's unrealistic, there is a related phenomena of diet-induced thermogenesis[1] known informally as “the meat sweats”. Some students connect this to feeling quite hungry after a cold swim in the pool (a similar effect). On a larger scale, talking about what's wrong with this estimate is useful. The main storage mechanism for storing food energy is fat tissue, which the calculation completely ignores. Infants are generally born with little fat, and an infant sleeping through the night often coincides with the baby growing enough fat to store enough kcals to make it through a night without waking up ravenously hungry. A related follow-up is that if a person is stranded in the wilderness, they should immediately start walking downstream (ie, towards civilization) as they likely won't be able to harvest an amount of kcals equivalent to what they already have stored on their hips and abdomen.[?] The contrast of bear hibernation [?] and songbirds constantly eating through the winter are related connections to investigate.

Section Biophysical Power: A more realistic question to follow up with relates to the average *power* given off by a person over a day. Again, assuming 3000kcal is burned over 24 hours, with useful information:  $1\text{kcal} \approx 4200\text{J}$  and  $1\text{J}/\text{s} = 1\text{W}$ .

$$\frac{3000\text{kcal}}{24\text{hours}} \frac{4200\text{J}}{1\text{kcal}} \frac{1\text{hour}}{3600\text{sec}} \approx 145\text{W} \quad (3)$$

Most students still remember 75Watt lightbulbs, but given the spread of LED lighting, “A person is two 75W light bulbs” will probably only make sense for a few more years. Desert or cole-weather camping, alone or with friends, and survival swimming are also examples for students to make sense of this answer. If you can take advantage of other people's waste body heat, you'll sleep more pleasantly.

Another connection to make is that of “brown fat” a sort of biological space heater that humans can develop in response to cold weather. For example, this tissue's mitochondria can burn lipids and carbohydrates in a useless proton pumping scheme, which produces metabolic heat [?]. Most common in rodents and infants, this mechanism can be stimulated by extended exposure to cold temperatures. The idea of a biological

space heater that takes a month to turn on and a month to turn off matches the lived experience of college students in Minnesota, who wear down jackets in  $4C^\circ$  weather in November, and beachwear in  $4C^\circ$  weather in March. Additionally, transplants to northern climates often take a few years to “get used to” the colder weather up north. It seems just as easy to say that transplants’ bodies take a few years to grow the brown fat that allows them to be comfortable with cold weather.

Power is the slope of energy-time METS Imagine that after eating a 600 *kcal* bacon maple long-john (donut), you decide to go for a hike to work off the Calories. Winona State is in a river valley bounded by 200m tall bluffs. How high up the bluff would you have to hike to burn off the donut? Useful information: human muscle is about 30% efficient and gravitational energy on Earth’s surface has a slope of about 10 *Joules/kg · m*.

Answer Energy bar charts

$$\frac{1}{3} \cdot 600kcal \cdot \frac{4200J}{1kcal} = 80kg \cdot 10 \frac{Joules}{kg \cdot m} \cdot height \quad (4)$$

$$height \approx 1000m \quad (5)$$

increase in yields since 1917 (graph)

1917 data

Grow your own food, possible?

Grow your own food, how far apart (urban life?)

### 3. Example: How big could Tenochtitlan have been?

1917 (A&M) USDA pamphlet

Corn for US - area

If Tenoch was 100k people, how much land area?

### 4. Example: Was the Irish Potato Famine a Natural Disaster?

### 5. Conclusion

#### Acknowledgments

The work was prompted in part by discussions with John Deming, Carl Ferkinhoff, and Sarah Taber.

#### Appendix A. Introductory Food Energy Questions

Planning to save money, one college student decides to go to an all-you-can-eat buffet each day at 11am. If he brings homework and stretches the meal out for a few hours he can get all 3000 *kcal*s with only one meal bill. Food is fuel for the human body. If his body burned all this food at once, how much warmer would he get? Useful information:

the student has a mass of 80kg and is made mostly of water. A Calorie heats 1 kg of water  $1^{\circ}C$ .

Answer

$$3000kcal = 80kg \cdot 1 \frac{kcal}{kg \cdot C^{\circ}} \cdot \Delta T \quad (A.1)$$

$$\Delta T \approx +37.5^{\circ} \quad (A.2)$$

Fat tissue serves a valuable purpose, brown fat, babies, songbirds

What power does the body give off in the more realistic case that the 3000kcal is burned over 24 hours? Useful information:  $1kcal \approx 4200J$  and  $1J/s = 1W$ .

$$\frac{3000kcal}{24hours} \frac{4200J}{1kcal} \frac{1hour}{3600sec} \approx 145W \quad (A.3)$$

Survival swimming, putting all the kids in one bed on a cold winter night.

## References

- [1] <https://www.bonappetit.com/story/meat-sweats> P. Trayhurn, THERMOGENESIS, Editor(s): Benjamin Caballero, Encyclopedia of Food Sciences and Nutrition (Second Edition), Academic Press, 2003, Pages 5762-5767, ISBN 9780122270550, <https://doi.org/10.1016/B0-12-227055-X/01188-3>. (<https://www.sciencedirect.com/science/article/pii/B012227055X011883>)
- [?] The wilderness river might be full of trout, but if they're 300kcal each, you'll have to catch, clean, and smoke 10 of them to store up a day's food. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/175154/nutrients>
- [?] Some sources claim that bear metabolism can vary between 4000 to and 20,000 kcal per day, <https://bear.org/5-stages-of-activity-and-hibernation/>, comically illustrated by the National Park Service at <https://www.nps.gov/katm/learn/fat-bear-week-2022.htm>.
- [2] A space heater <https://hms.harvard.edu/news/research-brown-fat-heats>  
Brown and Beige Fat: Molecular Parts of a Thermogenic Machine Paul Cohen<sup>1</sup> and Bruce M. Spiegelman<sup>2</sup> Diabetes 2015;64:2346–2351 — DOI: 10.2337/db15-0318  
<https://pubmed.ncbi.nlm.nih.gov/33846638/> Shamsi F, Piper M, Ho LL, Huang TL, Gupta A, Streets A, Lynes MD, Tseng YH. Vascular smooth muscle-derived Trpv1+ progenitors are a source of cold-induced thermogenic adipocytes. Nat Metab. 2021 Apr;3(4):485-495. doi: 10.1038/s42255-021-00373-z. Epub 2021 Apr 12. PMID: 33846638; PMCID: PMC8076094.  
<https://pubmed.ncbi.nlm.nih.gov/14715917/> Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. Physiol Rev. 2004 Jan;84(1):277-359. doi: 10.1152/physrev.00015.2003. PMID: 14715917.  
<https://pubmed.ncbi.nlm.nih.gov/6722594/> Himms-Hagen J. Nonshivering thermogenesis. Brain Res Bull. 1984 Feb;12(2):151-60. doi: 10.1016/0361-9230(84)90183-7. PMID: 6722594.  
<https://www.nih.gov/news-events/nih-research-matters/uncovering-origins-brown-fat>
- [3] for crop productivity
- [4]